Supplementary Table 1. Comparison of the available baseline characteristics between the included and excluded patients					
Variables	**Final cohort (n*)**		**Excluded patients (n)**		**p**
Age at first ^{90}Y SIRT, mean ± SD	79	63.1±9.2 years	24	63.7±10 years	0.783
BCLC stage, n (%)	79		19		
B		43 (54.4)		13 (68.4)	0.269
C		36 (45.6)		6 (31.6)	
Child-Pugh score, n (%)	73		20		
A		63 (79.7)		15 (75)	0.301
B		10 (12.7)		5 (25)	
Extrahepatic metastasis, n (%)	79	15 (19)	24	3 (13)	0.555
Laboratory values, median (min-max)					
Albumin, mean ± SD	94	3.7±0.5 g/dL	24	3.6±0.5 g/dL	0.487
Total bilirubin	94	0.8 (0.3-2.4) mg/dL	23	0.7 (0.3-2.8) mg/dL	0.853
ALT	94	32 (8-260) U/L	21	52 (14-102) U/L	0.099
AST	94	43 (13-248) U/L	21	65 (22-162) U/L	0.251
INR	81	1.1 (0.8-1.5)	20	1.1 (0.9-1.8)	0.371
Liver function index, median (min-max)					
ALBI score, mean ± SD	94	-2.36±0.49	23	-2.27±0.53	0.437

*: Laboratory values and index of liver function are presented on a per-treatment basis, as these parameters could vary between sequential/repetitive ^{90}Y SIRT sessions in the same patient
^{90}Y SIRT: Yttrium-90 selective internal radiation therapy, BCLC: Barcelona Clinic Liver Cancer, ALT: Alanine aminotransferase, AST: Aspartate aminotransferase, INR: International normalized ratio, ALBI: Albumin-bilirubin, SD: Standard deviation, min: Minimum, max: Maximum